

J. H. Choi · S. Y. Lee

Secretory and extracellular production of recombinant proteins using *Escherichia coli*

Received: 25 September 2003 / Revised: 25 December 2003 / Accepted: 30 December 2003 / Published online: 14 February 2004
© Springer-Verlag 2004

Abstract *Escherichia coli* is one of the most widely used hosts for the production of recombinant proteins. However, there are often problems in recovering substantial yields of correctly folded proteins. One approach to solve these problems is to have recombinant proteins secreted into the periplasmic space or culture medium. The secretory production of recombinant proteins has several advantages, such as simplicity of purification, avoidance of protease attack and N-terminal Met extension, and a better chance of correct protein folding. In addition to the well-established Sec system, the twin-arginine translocation (TAT) system has recently been employed for the efficient secretion of folded proteins. Various strategies for the extracellular production of recombinant proteins have also been developed. For the secretory production of complex proteins, periplasmic chaperones and protease can be manipulated to improve the yields of secreted proteins. This review discusses recent advances in secretory and extracellular production of recombinant proteins using *E. coli*.

Introduction

Escherichia coli has been the “workhorse” for the production of recombinant proteins as it is the best-characterized host with many available expression systems

J. H. Choi · S. Y. Lee
Metabolic and Biomolecular Engineering National Research Laboratory, Department of Chemical and Biomolecular Engineering and BioProcess Engineering Research Center, 373-1 Guseong-dong, Korea Advanced Institute of Science and Technology, 305-701 Yuseong-gu, Daejeon, Korea

S. Y. Lee (✉)
Department of Biosystems and Bioinformatics Research Center, Korea Advanced Institute of Science and Technology, 373-1 Guseong-dong, 305-701 Yuseong-gu, Daejeon, Korea
e-mail: leesy@kaist.ac.kr
Tel.: +42-869-3930
Fax: +42-869-8800

(Lee 1996; Makrides 1996). However, *E. coli* cannot produce some proteins containing complex disulfide bonds, or mammalian proteins that require post-translational modification for activity. Nevertheless, many recombinant proteins have been successfully produced using *E. coli*. Overexpressed proteins are often produced in the form of inclusion bodies, from which biologically active proteins can only be recovered by complicated and costly denaturation and refolding processes. Furthermore, the final yields of these soluble refolded proteins are usually very low, due mainly to protein aggregation resulting from interactions between the hydrophobic regions of the proteins.

A variety of techniques have been developed to solve these problems, including the use of different promoters to regulate the level of expression, the use of different host strains, co-expression of chaperones, reduction of culture temperature, and secretion of proteins into the periplasm or culture medium. In this review, we discuss strategies for secretory and extracellular production of recombinant proteins using *E. coli*.

Secretory production of recombinant proteins using *E. coli*

Secretory production of recombinant proteins provides several advantages compared to cytosolic production. For example, the N-terminal amino acid residue of the secreted product can be identical to the natural gene product after cleavage of the signal sequence by a specific signal peptidase. Also, there appears to be much less protease activity in the periplasmic space than in the cytoplasm. In addition, recombinant protein purification is simpler due to fewer contaminating proteins in the periplasm. Another advantage is that correct formation of disulfide bonds can be facilitated because the periplasmic space provides a more oxidative environment than the cytoplasm (Makrides 1996). Furthermore, it has recently been shown that secreted proteins can be applied in in-vivo activity assays as secretion gives the expressed protein (enzyme) greater

access to the substrate. Secreted proteins can also be used to screen protein libraries (Chen et al. 2001; Sroga and Dordick 2002).

In gram-negative bacteria, at least three different types of protein secretion systems have been described—type I, type II, and type III (Pugsley 1993), with type II being the most widely used. The type II system involves a two-step process in which a premature protein containing a signal sequence is exported to the periplasmic space using the Sec pathway and processed into a mature protein. Although heterologous proteins are often exported to the periplasm by the common Sec system, extracellular secretion of target proteins depends on the characteristics of signal sequences and proteins.

Generally, proteins found in the outer membrane or periplasmic space are synthesized in the cytoplasm as premature proteins. These premature proteins contain a short (15–30) specific amino acid sequence (signal sequence) that allows proteins to be exported outside the cytoplasm. A number of signal sequences have been used for efficient secretory production of recombinant proteins in *E. coli*, including PelB, OmpA, PhoA, endoxylanase, and StII. Although sequence diversity exists among these signal sequences, some common structural features have been identified. Table 1 shows the representative signal sequences that have been used for the secretory production of recombinant proteins in *E. coli*. Typical signal sequences are composed of a hydrophobic H-domain of about 10–20 amino acids that can be preceded by a short, positively charged N-domain of about two to ten amino acids (Fig. 1, Table 1). In general, signal sequences are rich in hydrophobic amino acids, such as alanine, valine, and leucine, a feature essential for secretion of the proteins into the periplasm of *E. coli* (Pugsley 1993). During transport of proteins out of the cytoplasm, the signal sequence is cleaved by signal peptidase to yield a mature protein product. The cleavage site (the C-domain) is usually less hydrophobic, contains a signal recognized by the signal peptidase, and conforms to the $-3, 1$ rule (Pugsley 1993)—that is, the residue at position -1 must have a small neutral side-chain, as is the case for alanine,

glycine, and serine, and this also holds true for the residue at position -3 . As can be seen from Table 1, alanine is most frequently found at the -1 and -3 positions, forming the so-called Ala-X-Ala box, which is recognized and cleaved by signal peptidase I. The secondary structure at the cleavage junction of preproteins also plays an important role in determining the cleavage site and protein processing (Pratap and Dikshit 1998). Thus, the selection of an optimal signal sequence is important for efficient secretory production of recombinant proteins. As can be seen from Table 2, the efficiency of protein secretion varies depending on the host strain, signal sequence, and the type of protein to be secreted. To date, there is no general rule in selecting a proper signal sequence for a given recombinant protein to guarantee its successful secretion. Several signal peptides, such as those listed in Table 1, must be examined in a trial-and-error type approach.

One of the advantages of secretory protein production is that the authentic N-terminal amino acid sequence without the Met extension can be obtained after cleavage by the signal peptidase. However, this can be achieved only when the gene of interest is correctly fused to the cleavage site. Choi et al. (2000) reported on the use of a new signal sequence cloned from the *Bacillus* sp. endoxylanase gene (Jeong et al. 1998) for the secretory production of recombinant proteins. Within the signal peptidase cleavage site (A-S-A) of the endoxylanase signal sequence, there is a *PstI* site that can be used for directly cloning the gene encoding the mature protein of interest. Upon cleavage by the signal peptidase, the mature protein can be produced without the need to change any amino acid sequence. Therefore, the endoxylanase signal sequence allows convenient cloning of genes encoding recombinant proteins for secretory production without changing either the sequence itself or the sequence of the mature protein. This is an important feature of the endoxylanase signal peptide as other signal sequences often cannot be used without changing the mature protein sequence. Using the endoxylanase signal sequence and the inducible *trc* promoter, *Bacillus* sp. endoxylanase and *E. coli* alkaline

Table 1 Representative signal sequences used for the secretory production of recombinant proteins in *Escherichia coli*. The signal sequence is composed of N-, H- and C-domains. The N-domains of signal sequences are shown in *bold* while the C-domains are *underlined*

Signal sequences	Amino acid sequences
PelB (pectate lyase B) from <i>Erwinia carotovora</i>	MK YLLPTAAAGLLLLAAQ PAMA
OmpA (outer-membrane protein A)	MK KTAIAIAVALAG FATVAQA
StII (heat-stable enterotoxin 2)	MK KNIAFLLASMFVFSIAT NAYA
Endoxylanase from <i>Bacillus</i> sp.	MF KFKKKFLVGLTAA FMSISMFSATASA
PhoA (alkaline phosphatase)	MK QSTIALALLPL LFTPVTKA
OmpF (outer-membrane protein F)	MM KRNILAVIVPALLV AGTANA
PhoE (outer-membrane pore protein E)	MK KSTLALVVMGIVAS SASVQA
MalE (maltose-binding protein)	MK IKTGARILALSALT TMMFSASALA
OmpC (outer-membrane protein C)	MK VKVLSSLVPALLV AGAANA
Lpp (murein lipoprotein)	MK ATKLVLGAVIL GSTLLAG
LamB (λ receptor protein)	MM ITLRKLP LAVAVAAGVMSAQAMA
OmpT (protease VII)	MR AKLLGIVL TTPIAISSFA
LTB (heat-labile enterotoxin subunit B)	MN KVKCYVL FTALLSSLYAHG




Fig. 1A–C Strategies for the secretory production of recombinant proteins in *Escherichia coli*. The Sec system, the twin-arginine translocation (TAT) system, and the strategies for enhancing secretory protein production using periplasmic chaperones and protease-negative mutants are shown. **A** The co-expression of periplasmic chaperones, such as disulfide-bond formation (Dsb) family proteins, SurA, FkpA, and Skp, can improve the efficiencies of secretory production and protein folding (Arie et al. 2001;

Bothmann and Pluckthun 1998, 2000; Jeong and Lee 2000; Kurokawa et al. 2001; Lazar and Kolter 1996; Qiu et al. 1998; Wulfiging and Rappouoli 1997; Zavialov et al. 2001). **B** Protease-negative mutant strains can improve secretory production of recombinant proteins by reducing proteolysis (Park et al. 1999; Wulfiging and Rappouoli 1997). **C** A novel TAT system can directly secrete the folded proteins (Angelini et al. 2001; Barrett et al. 2003; Santini et al. 2001; Thomas et al. 2001)

phosphatase could be secreted into the *E. coli* periplasm. Furthermore, human leptin and granulocyte-colony stimulating factor (G-CSF) were also efficiently secreted into the *E. coli* periplasm using this sequence (Jeong and Lee 2000, 2001; Yim et al. 2001).

More recently, a novel twin-arginine translocation (TAT) system was discovered. The TAT system is Sec independent and is capable of secreting folded proteins by employing a particular signal peptide containing a twin-arginine sequence. Green fluorescent protein (GFP) fused to the twin-arginine signal peptide of trimethylamine-*N*-oxide (TMAO) reductase (TorA), and alkaline phosphatase (Tap) from *Thermus thermophilus*, which also uses the TAT system, were successfully secreted into the periplasm of *E. coli* (Angelini et al. 2001; Barrett et al. 2003; Santini et al. 2001; Thomas et al. 2001). Since the TAT system allows secretion of proteins already folded in the *E. coli* cytosol, it may have advantages over the Sec system, particularly for those proteins that may be folded before they can reach the Sec machinery or that contain complex disulfide bonds.

Despite many successful examples, such as those described above, secretory production of heterologous

proteins in *E. coli* remains problematic. Obstacles include: (1) incomplete processing of signal sequences, (2) variable secretion efficiency depending on the characteristics of the proteins, (3) low or undetectable amounts of recombinant protein secretion, (4) formation of inclusion bodies in the cytosol and periplasm when using strong promoters, and (5) incorrect formation of disulfide bonds (Chung et al. 1998; Jeong and Lee 2000; Lucic et al. 1998; Pritchard et al. 1997; Wong et al. 2003). The first three problems have been solved using a trial-and-error type approach; different promoters, signal sequences, and host strains were examined under various culture conditions (e.g. temperature). The third problem might be due to periplasmic proteolysis rather than poor secretion machinery. In this case, a host strain deficient in periplasmic proteases can be used (see below). The fourth and fifth problems have been solved by manipulating periplasmic chaperones, as shown below. In addition, the use of the TAT system may be a good alternative to solve some of these problems.

Table 2 Representative secretory production of recombinant proteins in *E. coli*

Model proteins	Signal sequences	Hosts	Characteristics	References
Staphylokinase (sak)	OmpA	<i>E. coli</i> JM109	15 µg/ml	Lee et al. 1998
Human leptin	Endoxylanase	<i>E. coli</i> BL21(DE3)	41% of total proteins, DsbA coexpression	Jeong and Lee 2000
Staphylokinase (sak)	Native, PelB, LambB, MeIE, OmpA	<i>E. coli</i>	140 IU/ml (periplasm), 530 IU/ml (extracellular)	Pratap and Dikshit 1998
Green fluorescent protein (GFP)	TorA (TMAO reductase)	<i>E. coli</i> MC4100	TAT pathway	Barrett et al. 2003; Santini et al. 2001; Thomas et al. 2001
Human interleukin-2 receptor α-chain (hIL-2Rα)	StII	<i>E. coli</i> DH5α		Dracheva et al. 1995
<i>Arthrobacter levan</i> fructotransferase	LacZ derived secretion motif	<i>E. coli</i> JM109	2 g/l	Lee et al. 2001
Mutated heat-labile enterotoxin	Native	<i>E. coli</i> KS476 (<i>degP</i>)	DsbA coexpression, DegP knock-out strain	Wulfling and Rappouli 1997
Thermoalkaliphilic lipase	Native, OmpA	<i>E. coli</i> JM105	660000 U/ml	Rua et al. 1998
Cholera toxin, and B subunit	Heat-labile enterotoxin LTIIb	<i>E. coli</i> TX1	190 mg/ml	Jobling et al. 1997
Alkaline phosphatase	Endoxylanase	<i>E. coli</i> HB101	5.2 g/l	Choi et al. 2000
Human interleukin-1β-cafl fusion protein	CafI	<i>E. coli</i> JM105	Chaperone/usher pathway	Zavialov et al. 2001
Mouse endostatin	PhoA	<i>E. coli</i> DH5α	40 mg/l	Xu et al. 2002
Subtilisin E	Gene III	<i>E. coli</i> TOP10	Cell-based kinetic assay	Sroga and Dordick 2002
Penicillin G acylase	Native (IB)	<i>E. coli</i> DH5α		Sriubolmas et al. 1997
Human granulocyte colony- stimulating factor (GCSF)	Endoxylanase	<i>E. coli</i> BL21(DE3)	22% of total proteins	Jeong and Lee 2001; Yim et al. 2001
Protein A-β-lactamase	Protein A	<i>E. coli</i> HM114	112500 U/l	Park et al. 1999
Hirudin III	L-Asparaginase II	<i>E. coli</i> JM105	60 mg/l	Tan et al. 2002
Protein A-PhoA fusion protein	PelB	<i>E. coli</i> BL21(DE3)		Chowdhury et al. 1994
Penicillin acylase (PAC)	Native	<i>E. coli</i> HB101	DegP coexpression	Lin et al. 2001a
Human cytochrome P4501A1	PhoA	<i>E. coli</i> TB1	4500 nmol/l	Kaderbhai et al. 2000
<i>Manduca</i> diuresin	OmpA	<i>E. coli</i> JM101	Proteolysis	Wong et al. 2003
scFv antibody	OmpA	<i>E. coli</i> JM109	Bacteria L form	Rippmann et al. 1998
Pullulanase (PulA)	Naive	<i>E. coli</i> K12	dsbA effect	Pugsley et al. 2001
Human proinsulin	SpA	<i>E. coli</i> AF1000	4.6 mg/l, <i>uspA</i> and <i>uspB</i> promoter	Mergulhao et al. 2003
20-kDa human growth hormone	<i>npr</i> (neutral protease gene), ompA	<i>E. coli</i> W3110	76 mg/l, glutathione reductase coexpression	Uchida et al. 1997
Human granulocyte-colony stimulating factor (hGCSF)	PelB	<i>E. coli</i> BL21(DE3)	Not secreted	Chung et al. 1998
Human, mouse leptin	OmpA	<i>E. coli</i> MC1061	1.46 mg/ml	Guisez et al. 1998
Peptide-N-glycosidase F (PNGase F)	OmpA	<i>E. coli</i> BL21(DE3)	8 mg/l	Loo et al. 2002

Table 2 (continued)

Model proteins	Signal sequences	Hosts	Characteristics	References
<i>Thermus thermophilus</i> alkaline phosphatase (Tap)	Native	<i>E. coli</i> CC118	TAT pathway	Angelini et al. 2001
β hCG (chorionic gonadotropin)	LTB (heat labile enterotoxin)	<i>E. coli</i> DH5 α		Pillai et al. 1996
Human proinsulin	DabA fusion	<i>E. coli</i> RB791	9.2 mg/g, Arg and ethanol addition	Winter et al. 2000
Alkaline phosphatase	Degenerated PelB	<i>E. coli</i> W3110		Le Calvez et al. 1996
Tn <i>phoA</i>	SpaA fusion at different site	<i>E. coli</i> CC202		Holt and Raju 2000
Human nerve growth factor β	OmpT	<i>E. coli</i> JM109	Dsb protein coexpression	Kurokawa et al. 2001
Human tissue plasminogen activator	StII	<i>E. coli</i> SF110	Dsb protein coexpression	Qiu et al. 1998

Enhancement of secretion efficiency by manipulating periplasmic chaperones

Many proteins contain disulfide bonds that need to be correctly formed to be functional. Some proteins containing disulfide bonds can be produced in active forms in the periplasm or extracellular medium using the secretion system. However, production of large and complex recombinant proteins in the *E. coli* periplasm can be limited by low secretion levels and folding problems, leading to periplasmic inclusion bodies. Many periplasmic chaperones have been characterized, and subsequently used for the efficient secretion of recombinant proteins. These proteins assist in correct folding of secreted proteins and prevent the formation of periplasmic inclusion bodies. Strategies for enhancing secretion efficiency of recombinant proteins in *E. coli* are presented in Fig. 1.

Large and complex proteins from mammalian cells frequently contain disulfide bonds which contribute to their stability and, in many cases, are essential for their catalytic activities. However, the *E. coli* cytosol is a rather reduced environment, and thus disulfide bonds are not normally formed. The enzymes that catalyze disulfide bond formation play key roles in folding many secreted proteins. For example, the Dsb (disulfide-bond formation) family of proteins catalyzes both the formation of new disulfide bonds and the rearrangement of existing ones. Dsb proteins contain one or more highly conserved thioredoxin-like motifs (C-X-X-C) which are important for disulfide oxidoreductase activity. DsbA and DsbB are oxidoreductases that allow the formation of disulfide bonds (Fig. 1). Subsequent rearrangement of the newly formed disulfide bonds is sometimes necessary since they can be formed among incorrectly paired cysteines, trapping substrate proteins in a misfolded conformation. Normally, misfolded proteins in the periplasm do not readily accumulate, being rapidly degraded by DegP protease. Disulfide-bond rearrangement is catalyzed by two periplasmic disulfide bond isomerases, DsbC and DsbD (Fig. 1). Recent studies revealed that the overexpression of Dsb proteins increased secretion efficiency, folding, and the solubility of recombinant proteins in the periplasmic space (Jeong and Lee 2000; Kurokawa et al. 2001; Qiu et al. 1998; Wulping and Rappouli 1997).

SurA, FkpA, and Skp are another set of folding catalysts and chaperones in the periplasmic space (Bothmann and Pluckthun 2000; Missiakas et al. 1996). SurA, which was identified in *E. coli* during starvation survival, shares sequence similarity with parvulin, a cytoplasmic peptidyl prolyl isomerase (PPI) in *E. coli*. Lazar and Kolter (1996) demonstrated that SurA assists in periplasmic folding of three outer-membrane proteins (OmpA, OmpF, and LamB) and of some other secreted proteins. SurA promotes folding of several otherwise unstable proteins (e.g. Protein A- β -lactamase hybrid protein) and proteins prone to aggregation. FkpA, which is a heat-shock periplasmic peptidylprolyl *cis/trans* isomerase, was shown to suppress the formation of modified maltose-binding protein (MalE31) inclusion bodies and to enhance

periplasmic expression and folding efficiency of functional antibody fragments in *E. coli* (Arie et al. 2001; Bothmann and Pluckthun 2000). Skp is also a periplasmic chaperone and it appears to improve phage-antibody display and periplasmic expression (Bothmann and Pluckthun 1998).

Sometimes, efficient protein secretion is hindered due to the degradation of target proteins by cell-envelope proteases, such as DegP, OmpT, protease III, and Tsp. Strategies to minimize proteolysis include reducing cultivation temperatures and using protease-negative mutant strains (Park et al. 1999; Wulfiging and Rappuoli 1997).

Zavialov et al. (2001) demonstrated another novel approach using the chaperone/usher pathway for the secretion of human interleukin-1 β and F1 antigen (Caf1) fusion protein into the periplasmic space. In this system, the Caf1M chaperone assisted periplasmic folding and enhanced chimeric protein solubilization.

Extracellular production of recombinant proteins

Extracellular production of recombinant proteins has several advantages over secretion into the periplasm (Shokri et al. 2003). Extracellular production does not require outer-membrane disruption to recover target proteins, and, therefore, it avoids intracellular proteolysis by periplasmic proteases and allows continuous production of recombinant proteins. Based on these advantages, various strategies have been developed in *E. coli* for the extracellular production of recombinant proteins (Fig. 2, Table 3).

A number of methods have been applied to promote extracellular secretion of recombinant proteins from *E. coli*. These include the use of biochemicals, physical methods (osmotic shock, freezing and thawing), lysozyme treatment, and chloroform shock. However, these methods can be applied only after harvesting cells. *E. coli* normally does not secrete proteins extracellularly except for a few classes of proteins such as toxins and hemolysin. Secreted proteins can leak from the periplasmic space into the




Fig. 2A–D Strategies for the extracellular production of recombinant proteins by *E. coli*. **A** Recombinant proteins can be excreted into the culture medium by treating cells with various agents or by using L-form cells (Jang et al. 1999; Kaderbhai et al. 1997; Kujau et al. 1998; Yang et al. 1998). **B** Recombinant protein fused to outer-membrane protein F (OmpF) of *E. coli* can be excreted into the culture medium (Jeong and Lee 2002; Nagahari et al. 1985). **C** Proteins secreted into the *E. coli* periplasm can also be released into the culture medium by co-expression of *kil*, *out* genes, the gene

encoding the third topological domain of the transmembrane protein TolA (TolAIII), or the bacteriocin-release protein gene (Fu et al. 2003; Kleist et al. 2003; Lin et al. 2001b; Miksch et al. 1997; 2002; Robbens et al. 1995; van der Wal et al. 1995, 1998; Wan and Baneyx 1998; Zhou et al. 1999). **D** The target protein fused to the C-terminal hemolysin secretion signal can be directly excreted into the culture medium through the hemolysin transport system (Fernandez et al. 2000; Li et al. 2002)

Table 3 Representative extracellular production of recombinant proteins in *E. coli*

Model proteins	Signal sequences	Hosts	Characteristics	References
Levansucrase, β -lactamase	Native	<i>E. coli</i> DH5 α	5% of total proteins, glycine supplement	Jang et al. 1999
Winter flounder antifreeze	OmpA	<i>E. coli</i> JM105	16 mg/l	Tong et al. 2000
Human β -endorphin	OmpF hybrid	<i>E. coli</i> BL21(DE3)	Hybrid proteins in medium	Jeong and Lee 2002
Exoglucanase	OmpA	<i>E. coli</i> JM101	143 U/ml	Lam et al. 1997
Human epidermal growth factor (hEGF)	OmpA	<i>E. coli</i> JM101	325 mg/l	Sivakesava et al. 1999
Taq I restriction endonuclease	MBP hybrid	<i>E. coli</i> XL1-Blue	30000 U/l	Toksoy et al. 2002
Pectate lyase	PelB	<i>E. coli</i> BL21(DE3)	2200 U/ml	Matsumoto et al. 2002
Fungal ribotoxin α -sarcin	OmpA	<i>E. coli</i> BL21(DE3)	3 μ g/ml	Rathore et al. 1997
Fimbrial molecular chaperone FaeE	Native	<i>E. coli</i> BL21(DE3)	21 mg/l, bacteriocin-release protein	van der Wal et al. 1995
Fimbrial molecular chaperone FaeE	Native	<i>E. coli</i> C600	Modified bacteriocin-release protein	van der Wal et al. 1998
Penicillin acylase	Native	<i>E. coli</i> C600	Bacteriocin-release protein	Lin et al. 2001b
β -Glucanase	Native	<i>E. coli</i> HB101, MD Δ P7	Bacteriocin-release protein	Lin et al. 2001b
scFv Antibody	OmpA,	<i>E. coli</i> JM109	150 U/ml, <i>kil</i> coexpression	Miksch et al. 1997
Phytase	Native	<i>E. coli</i> K802	170-fold increased, glycine, Triton X-100	Yang et al. 1998
Phytase	OmpA,	<i>E. coli</i> K802	Hemolysin system	Fernandez et al. 2000
<i>Erwinia chrysanthemi</i> endoglucanase	Native	<i>E. coli</i> HB2151	50 U/ml, <i>kil</i> coexpression	Miksch et al. 2002
Mini-antibodies McPC603scFvDhlx	Native	<i>E. coli</i> BL21(DE3)	120 U/ml, <i>kil</i> coexpression, HCDC	Kleist et al. 2003
Insulin-like growth factor binding protein-2 (IGFBP-2)	Native	<i>E. coli</i> BL21(DE3)	4-6% of total proteins, <i>out</i> coexpression	Zhou et al. 1999
Murine interleukin-2 (mIL2)	OmpA	<i>E. coli</i> B	10 mg/l, bacterial L-form	Kujau et al. 1998
Heat stable alkaline protease	PelB	<i>E. coli</i> RV308	Gene III product fusion	Lucic et al. 1998
TEM- β -lactamase	OmpA	<i>E. coli</i> BL21	16 mg/l, <i>kil</i> gene	Robbens et al. 1995
Human C-reactive protein (CRP)	Native	<i>E. coli</i> K12 RP1 Δ M15	1500 U/ml, bacteriocin-release protein	Fu et al. 2003
Human interleukin 6	OmpA	<i>E. coli</i> XL1-Blue	TolAIII coexpression	Wan et al. 1998
	PhoA (ALP)	<i>E. coli</i> BL21(DE3)	<i>kil</i> coexpression	Tanaka et al. 2002
	HlyA	<i>E. coli</i> JM109	70.4 ng/ml, hemolysin system	Li et al. 2002

culture medium possibly due to an increased permeability of the cell membrane during a lengthy incubation period. Small proteins secreted into the periplasm are frequently released into the culture medium (Tong et al. 2000). In general, movement of recombinant proteins from the periplasm to the culture medium is the result of compromising the integrity of the outer membrane. However, care must be exercised during such recombinant protein production so as not to compromise cellular integrity, which often causes cell death. Interestingly, glycine or Triton X-100 supplemented to the medium retarded formation of inclusion bodies in the periplasm and increased the extracellular production efficiency of recombinant proteins (Jang et al. 1999; Kaderbhai et al. 1997; Yang et al. 1998). Glycine has been found to induce morphological changes, such as an enlarged spheroidal morphology in *E. coli*, as it is incorporated into peptidoglycan. Glycine supplementation may slightly disrupt peptidoglycan cross-linkages and cell membrane integrity. Yang et al. (1998) reported that adding 2% (w/v) glycine dramatically increased extracellular production of sFV/TNF- α and β -glucosidase.

Another method of extracellular protein production involves fusing the product to a carrier protein that is normally secreted into the medium (e.g. hemolysin), or to a protein expressed on the outer membrane (e.g. OmpF). For example, human β -endorphin could be secreted into the culture medium when fused to OmpF (Jeong and Lee 2002; Nagahari et al. 1985). Recently, a method of releasing active scFv antibody and human interleukin-6 into the culture medium using the hemolysin secretion pathway was reported (Fernandez et al. 2000; Li et al. 2002). The hemolysin transport system (Hly) is a type-I secretory apparatus that forms a protein channel between the inner and outer membranes of *E. coli*. Hemolysin toxin (HlyA) is secreted by direct passage of the HlyA polypeptide from the cytoplasm to the extracellular medium using the hemolysin transport system (Fig. 2). For extracellular production using the hemolysin secretion pathway, the target protein is fused to the C-terminal hemolysin secretion signal. The Hly system appears to be an attractive candidate for the extracellular production of recombinant proteins.

Proteins secreted into the *E. coli* periplasm can also be released into the culture medium by co-expression of *kil* (Kleist et al. 2003; Miksch et al. 1997; 2002; Robbins et al. 1995) or the gene coding for the third topological domain of the transmembrane protein TolA (TolAIII) (Wan and Baneyx 1998). Zhou et al. (1999) reported extracellular production of the *Erwinia chrysanthemii*-doglucanase by employing the *out* genes from *E. chrysanthemi* EC16, which are responsible for the efficient extracellular secretion of pectic enzymes. Co-expression of the *out* genes increased production of active endoglucanase and released enzymes equivalent to over half of the total activity into the extracellular medium.

Another approach to the extracellular production of target proteins uses L-form cells, wall-less, or wall-deficient cells (Kujau et al. 1998; Rippmann et al.

1998). Recently, Kujau et al. (1998) demonstrated that L-form *E. coli* cells were capable of secreting into the culture medium a recombinant antibody fragment (single-chain phosphorylcholine-binding scFv from human McPC603) fused to the OmpA signal sequence under the control of the *lac* promoter. A correctly folded and dimerized mini-antibody was secreted directly, and remained stable in the culture medium.

Bacteriocin release protein (BRP) can also be used in the extracellular production of recombinant proteins in *E. coli*. BRP is a 28-amino-acid lipoprotein that activates detergent-resistant phospholipase A, resulting in the formation of permeable zones in the cell envelope through which proteins can pass into the culture medium (Fu et al. 2003; Lin et al. 2001b; van der Wal et al. 1995). However, co-expression of the BRP gene can damage the cell envelope and cause release of other cellular proteins. Recently, van der Wal et al. (1998) reported that a modified BRP gene (Lpp-BRP) could be used for the extracellular production of K88 fimbrial molecular chaperone FaeE without growth inhibition, lysis, or contaminating proteins.

Conclusions

Recent advances in our understanding of the protein secretory machinery and mechanism in *E. coli* have led to the development of various strategies to enhance secretory production of recombinant proteins. However, despite the successful development of various recombinant protein secretion systems, several problems remain to be solved. First, many large and complex proteins of eukaryotic origin are not efficiently secreted. Second, without trial-and-error, it is somewhat difficult to select a proper host-vector system and a signal sequence for the secretion of a desired protein. Third, the high-cell-density culture techniques for the secretory production of recombinant proteins are less well developed than those for cytosolic production. The first and second problems will be solved as our understanding of protein secretion pathways, folding mechanisms, periplasmic chaperone function, and signal sequences advances further. An obvious alternative solution to the first problem is simply not to adhere to the *E. coli* expression system, and instead use other organisms, including mammalian cells, as is currently practiced for a number of mammalian proteins. The third problem is that less research has been devoted to secretory production than to cytosolic expression, and therefore, can be solved by more research into the former. *E. coli* has been successfully used for both industrial- and laboratory-scale cytosolic production of recombinant proteins. Similar success with secretory and extracellular production of recombinant proteins using *E. coli* will likely follow.

Acknowledgements This review was supported by the Korean Systems Biology Research Grant (M10309020000-03B5002-00000) from the Ministry of Science and Technology. Support from IBM through the IBM-SUR program is greatly appreciated.

References

- Angelini S, Moreno R, Gouffi K, Santini C, Yamagishi A, Berenguer J, Wu L (2001) Export of *Thermus thermophilus* alkaline phosphatase via the twin-arginine translocation pathway in *Escherichia coli*. FEBS Lett 506:103-107
- Arie J, Sassoon N, Betton J (2001) Chaperone function of FkpA, a heat shock prolyl isomerase, in the periplasm of *Escherichia coli*. Mol Microbiol 39:199-210
- Barrett CML, Ray N, Thomas JD, Robinson C, Bolhuis A (2003) Quantitative export of a reporter protein, GFP, by the twin-arginine translocation pathway in *Escherichia coli*. Biochem Biophys Res Comm 304:279-284
- Bothmann H, Pluckthun A (1998) Selection for a periplasmic factor improving phage display and periplasmic expression. Nat Biotechnol 16:376-380
- Bothmann H, Pluckthun A (2000) The periplasmic *Escherichia coli* peptidylprolyl *cis,trans*-isomerase FkpA. J Biol Chem 275:17100-17105
- Chen G, Hayhurst A, Thomas JG, Garvey BR, Iverson BL, Georgiou G (2001) Isolation of high-affinity ligand-binding proteins by periplasmic expression with cytometric screening (PECS). Nat Biotechnol 19:537-542
- Choi JH, Jeong KJ, Kim SC, Lee SY (2000) Efficient secretory production of alkaline phosphatase by high cell density culture of recombinant *Escherichia coli* using the *Bacillus* sp. Endoxylanase signal sequence. Appl Microbiol Biotechnol 53:640-645
- Chowdhury PS, Kushwaha A, Abrol S, Chaudhary VK (1994) An expression system for secretion and purification of a genetically engineered thermostable chimera of protein A and alkaline phosphatase. Protein Expr Purif 5:89-95
- Chung BH, Sohn M, Oh S, Park U, Poo H, Kim BS, Yu MJ, Lee SK (1998) Overproduction of human granulocyte-colony stimulating factor fused to the PelB signal peptide in *Escherichia coli*. J Ferm Bioeng 85:443-446
- Dracheva S, Palermo RE, Powers GD, Waugh DS (1995) Expression of soluble human interleukin-2 receptor α -chain in *Escherichia coli*. Protein Expr Purif 6:737-747
- Fernandez LA, Sola I, Enjuanes L, De Lorenzo V (2000) Specific secretion of active single-chain Fv antibodies into the supernatants of *Escherichia coli* cultures by use of the hemolysin system. Appl Environ Microbiol 66:5024-5029
- Fu Z, Hamid SBA, Razak CAN, Basri M, Salleh AB, Rahman RNZA (2003) Secretory expression in *Escherichia coli* and single-step purification of a heat-stable alkaline protease. Protein Expr Purif 28:63-68
- Guisez Y, Fache I, Campfield LA, Smith FJ, Farid A, Plaetinck G, Van der Heyden J, Tavernier J, Fiers W, Burn P, Devos R (1998) Efficient secretion of biologically active recombinant OB protein (leptin) in *Escherichia coli*, purification from the periplasm and characterization. Protein Expr Purif 12:249-258
- Holt RG, Raju L (2000) Signal sequence and alanine-rich region of streptococcal protein antigen A of *Streptococcus sobrinus* can direct localization of alkaline phosphatase to the periplasm of *Escherichia coli*. FEMS Microbiol Lett 184:17-21
- Jang KH, Seo KB, Song KB, Kim CH, Rhee SK (1999) Extracellular secretion of levansucrase from *Zymomonas mobilis* in *Escherichia coli*. Bioproc Eng 21:453-458
- Jeong KJ, Lee PC, Park IY, Kim MS, Kim SC (1998) Molecular cloning and characterization of an endoxylanase gene of *Bacillus* sp. in *Escherichia coli*. Enzyme Microb Technol 22(7):599-605
- Jeong KJ, Lee SY (2000) Secretory production of human leptin in *Escherichia coli*. Biotechnol Bioeng 67:398-407
- Jeong KJ, Lee SY (2001) Secretory production of human granulocyte colony-stimulating factor in *Escherichia coli*. Protein Expr Purif 23(2):311-8
- Jeong KJ, Lee SY (2002) Excretory production of human β -endorphin into culture medium by using outer membrane protein F as a fusion partner in recombinant *Escherichia coli*. Appl Environ Microbiol 68:4979-4985
- Jobling MG, Palmer LM, Erbe JL, Holmes RK (1997) Construction and characterization of versatile cloning vectors for efficient delivery of native foreign proteins to the periplasm of *Escherichia coli*. Plasmid 38(3):158-73
- Kaderbhai MA, Ugochukwu CC, Lamb DC, Kelly S (2000) Targeting of active human cytochrome P4501A1 (CYP1A1) to the periplasmic space of *Escherichia coli*. Biochem Biophys Res Comm 279:803-807
- Kaderbhai N, Karim A, Hankey W, Jenkins G, Venning J, Kaderbhai MA (1997) Glycine-induced extracellular secretion of a recombinant cytochrome expressed in *Escherichia coli*. Biotechnol Appl Biochem 25:53-61
- Kleist S, Miksch G, Hitzmann B, Arndt M, Friehs K, Flaschel E (2003) Optimization of the extracellular production of a bacterial phytase with *Escherichia coli* by using different fed-batch fermentation strategies. Appl Microbiol Biotechnol 61:456-462
- Kujau MJ, Hoischen C, Riesenberger D, Gumpert J (1998) Expression and secretion of functional miniantibodies McPC603scFvDhx in cell-wall-less L-form strains of *Proteus mirabilis* and *Escherichia coli*: A comparison of the synthesis capacities of L-form strains with an *E. coli* producer strain. Appl Microbiol Biotechnol 49:51-58
- Kurokawa Y, Yanagi H, Yura T (2001) Overproduction of bacterial protein disulfide isomerase (DsbC) and its modulator (DsbD) markedly enhances periplasmic production of human nerve growth factor in *Escherichia coli*. J Biol Chem 276:14393-14399
- Lam T, Wong RSC, Wong WR (1997) Enhancement of extracellular production of a *Cellulomonas fimi* exoglucanase in *Escherichia coli* by the reduction of promoter strength. Enzyme Microb Technol 20:482-488
- Lazar SW, Kolter R (1996) SurA assists the folding of *Escherichia coli* outer membrane proteins. J Bacteriol 178:1770-1773
- Le Calvez H, Green JM, Baty D (1996) Increased efficiency of alkaline phosphatase production levels in *Escherichia coli* using a degenerate PelB signal sequence. Gene 170:51-55
- Lee J, Saraswat V, Koh I, Song KB, Park YH, Rhee SK (2001) Secretory production of *Arthrobacter* levan fructotransferase from recombinant *Escherichia coli*. FEMS Microbiol Lett 195(2):127-132
- Lee SJ, Kim IC, Kim DM, Bae KH, Byun SM (1998) High level secretion of recombinant staphylokinase into periplasm of *Escherichia coli*. Biotechnol Lett 20:113-116
- Lee SY (1996) High cell density cultivation of *Escherichia coli*. Trends Biotechnol 14:98-105
- Li Y, Chen CX, von Specht B, Hahn HP (2002) Cloning and hemolysin-mediated secretory expression of a codon-optimized synthetic human interleukin-6 gene in *Escherichia coli*. Gene 25:437-447
- Lin W, Huang S, Chou CP (2001a) DegP-coexpression minimizes inclusion-body formation upon overproduction of recombinant penicillin acylase in *Escherichia coli*. Biotechnol Bioeng 73:484-492
- Lin W, Huang S, Chou CP (2001b) High-level extracellular production of penicillin acylase by genetic engineering of *Escherichia coli*. J Chem Technol Biotechnol 76:1030-1037
- Loo T, Patchett ML, Norris GE, Lott JS (2002) Using secretion to solve a solubility problem: high-yield expression in *Escherichia coli* and purification of the bacterial glycoamidase PNGase F. Protein Expr Purif 24:90-98
- Lucic MR, Forbes BE, Grosvenor SE, Carr JM, Wallace JC, Forsberg G (1998) Secretion in *Escherichia coli* and phage-display of recombinant insulin-like growth factor binding protein-2. J Biotechnol J Biotechnol 61:95-108

- Makrides SC (1996) Strategies for achieving high-level expression of genes in *Escherichia coli*. *Microbiol Rev* 60:512–538
- Matsumoto T, Katsura D, Kondo A, Fukuda H (2002) Efficient secretory overexpression of *Bacillus subtilis* pectate lyase in *Escherichia coli* and single-step purification. *Biochem Eng J* 12(3):175–179
- Mergulhao FJM, Monteiro GA, Larsson G, Sanden AM, Farewell A, Nystrom T, Cabral JMS, Taipa MA (2003) Medium and copy number effects on the secretion of human proinsulin in *Escherichia coli* using the universal stress promoters *uspA* and *uspB*. *Appl Microbiol Biotechnol* 61:495–501
- Miksch G, Neitzel R, Fiedler E, Friehs K, Flaschel E (1997) Extracellular production of a hybrid β -glucanase from *Bacillus* by *Escherichia coli* under different cultivation conditions in shaking cultures and bioreactors. *Appl Microbiol Biotechnol* 47:120–126
- Miksch G, Kleist S, Friehs K, Flaschel E (2002) Overexpression of the phytase from *Escherichia coli* and its extracellular production in bioreactors. *Appl Microbiol Biotechnol* 56:685–694
- Missiakas D, Betton JM, Raina S (1996) New components of protein folding in extracytoplasmic compartments of *Escherichia coli* SurA, FkpA and Skp/OmpH. *Mol Microbiol* 21:871–884.
- Nagahari K, Kanaya S, Munakata D, Aoyagi Y, Mizushima S (1985) Secretion into the culture medium of a foreign gene product from *Escherichia coli*: use of the *ompF* gene for secretion of human β -endorphin. *EMBO J* 16:3589–3592
- Park SJ, Georgiou G, Lee SY (1999) Secretory production of recombinant protein by a high cell density culture of a protease negative mutant *Escherichia coli*. *Biotechnol Prog* 15:164–167
- Pillai D, Dixit A, Krishnan T, Garg LC (1996) Production of the beta-subunit of human chorionic gonadotropin in *Escherichia coli* and its export mediated by the heat-labile enterotoxin chain-B signal sequence. *Gene* 173(2):271–274
- Pratap J, Dikshit KL (1998) Efficient of signal peptide changes on the extracellular processing of streptokinase form *Escherichia coli* requirement for secondary structure at the cleavage junction. *Mol Gen Genet* 258:326–333
- Pritchard MP, Ossetian R, Li DN, Henderson CJ, Burchell B, Wolf CR, Friedberg T (1997) A general strategy for the expression of recombinant human cytochrome P450 s in *Escherichia coli* using bacterial signal peptides: Expression of CYP3A4, CYP2A6, and CYP2E1. *Arch Biochem Biophys* 345:342–354
- Pugsley AP (1993) The complete general secretory pathway in gram-negative bacteria. *Microbiol Rev* 57:50–108
- Pugsley AP, Bayan N, Sauvonnet N (2001) Disulfide bonds formation in secretion component PulK provides a possible explanation for the role of DsbA in pullulanase secretion. *J Bacteriol* 183:1312–1319
- Qui J, Swartz JR, Georgiou G (1998) Expression of active human tissue-type plasminogen activator in *Escherichia coli*. *Appl Environ Microbiol* 64:4891–4896
- Rathore D, Nayak SK, Batra JK (1997) Overproduction of fungal ribotoxin α -sarcin in *Escherichia coli*: generation of an active immunotoxin. *Gene* 190:31–35
- Rippmann JF, Klein M, Hoischen C, Brocks B, Rettig WJ, Gumpert J, Pfizenmaier K, Mattes R, Moosmayer D (1998) Prokaryotic expression of single-chain variable-fragment (scFv) antibodies: secretion in L-form cells of *Proteus mirabilis* leads to active product and overcomes the limitations of periplasmic expression in *Escherichia coli*. *Appl Environ Microbiol* 64(12):4862–4869.
- Robbens J, Raeymaekers A, Steidler L, Fiers W, Remaut E (1995) Production of soluble and active recombinant murine interleukin-2 in *Escherichia coli*: high level expression, *Kil*-induced release, and purification. *Protein Expr Purif* 6(4):481–486
- Rua ML, Atomi H, Schmidt-Dannert C, Schmid RD (1998) High-level expression of the thermoalkalophilic lipase from *Bacillus thermocatenulatus* in *Escherichia coli*. *Appl Microbiol Biotechnol* 49:405–410
- Santini C, Bernadac A, Zhang M, Chanal A, Ize B, Blanco C, Wu L (2001) Translocation of jellyfish green fluorescent protein via the Tat system of *Escherichia coli* and change of its periplasmic localization in response to osmotic up-shock. *J Biol Chem* 276:8159–8164
- Shokri A, Saden AM and Larsson G (2003) Cell and process design for targeting of recombinant protein into the culture medium of *Escherichia coli*. *Appl Microbiol Biotechnol* 60:654–664
- Sivakesava S, Xu ZN, Chen YH, Hackett J, Huang RC, Lam E, Lam TL, Siu KL, Wong RSC, Wong WKR (1999) Production of excreted human epidermal growth factor (hEGF) by an efficient recombinant *Escherichia coli* system. *Process Biochem* 34:893–900
- Sriubolmas N, Panbangred W, Sriuiratana S, Meevootisom V (1997) Localization and characterization of inclusion bodies in recombinant *Escherichia coli* cells overproducing penicillin G acylase. *Appl Microbiol Biotechnol* 47:373–378
- Sroga GE, Dordick J (2002) A strategy for in vivo screening of subtilisin E reaction specificity in *E. coli* periplasm. *Biotechnol Bioeng* 78:761–769
- Tan S, Wu W, Liu J, Kong Y, Pu Y, Yuan R (2002) Efficient expression and secretion of recombinant hirudin III in *E. coli* using the L-asparaginase II signal sequence. *Protein Expr Purif* 25:430–436
- Tanaka T, Horio T, Matuo Y (2002) Secretory production of recombinant human C-reactive protein in *Escherichia coli*, capable of binding with phosphorylcholine, and its characterization. *Biochem Biophys Res Commun*, 295(1):163–6
- Thomas JD, Daniel RA, Errington J, Robinson C (2001) Export of active green fluorescent protein to the periplasm by the twin-arginine translocase (Tat) pathway in *Escherichia coli*. *Mol Microbiol* 39:47–53
- Toksoy E, Onsan ZI, Kirdar B (2002) High-level production of *TaqI* restriction endonuclease by three different expression systems in *Escherichia coli* cells using the T7 phage promoter. *Appl Microbiol Biotechnol* 59:239–245
- Tong L, Lin Q, Wong WKR, Ali A, Lim D, Sung WL, Hew CL, Yang DSC (2000) Extracellular expression, purification, and characterization of a winter flounder antifreeze polypeptide form *Escherichia coli*. *Protein Expr Purif* 18:175–181
- Uchida H, Naito N, Asada N, Wada M, Ikeda M, Kobayashi H, Asanagi M, Mori K, Fujita Y, Konda K, Kusuhara N, Kamioka T, Nakashima K, Honjo M (1997) Secretion of authentic 20-kDa human growth hormone (20 K hGH) in *Escherichia coli* and properties of the purified product. *J Biotechnol* 55:101–112
- Van der Wal FJ, ten Hagen-Jounman CM, Oudega B, Luirink J (1995) Optimization of bacteriocin-release-protein-induced protein release by *Escherichia coli*: extracellular production of the periplasmic molecular chaperone FaeE. *Appl Microbiol Biotechnol* 44:459–465
- Van der Wal FJ, Koningstein G, ten Hagen CM, Oudega B, Luirink J (1998) Optimization of bacteriocin release protein (BRP)-mediated protein release by *Escherichia coli*: random mutagenesis of the pCloDF13-Derived BRP gene to uncouple lethality and quasi-lysis from protein release. *Appl Environ Microbiol* 64:392–398
- Wan EW, Baneyx F (1998) TolAIII co-expression facilitates the recovery of periplasmic recombinant proteins into the growth medium of *Escherichia coli*. *Protein Expr Purif* 14:13–22
- Winter J, Neubauer P, Glockshuber R, Rudolph R (2000) Increased production of human proinsulin in the periplasmic space of *Escherichia coli* by fusion to DsbA. *J Biotechnol* 84(2):175–185
- Wong WR, Ali AB, Ma MC (2003) Cloning, expression, and characterization of diuretic hormone *Manduca* diuresin from *Manduca sexta* in *Escherichia coli*. *Protein Expr Purif* 29:51–57
- Wulfling C, Rappuoli R (1997) Efficient production of heat-labile endotoxin mutant proteins by overexpression of *dabA* in *adegP*-deficient *Escherichia coli* strain. *Arch Microbiol* 167:280–283

- Xu R, Du P, Fan JJ, Zhang Q, Li TP, Gan RB (2002) High-level expression and secretion of recombinant mouse endostatin by *Escherichia coli*. *Protein Expr Purif* 24:453-459
- Yang J, Moyana T, Mackenzie S, Xia Q and Xiang J (1998) One hundred Seventy-fold increase in excretion of an FV frangmant tumor necrosis factor alpha fusion protein (SFV/TNF- α) from *Escherichia coli* caused by the synergistic effects of glycine and triton X-100. *Appl Environ Microbiol* 64:2669-2874
- Yim SC, Jeong KJ, Chang HN, Lee SY (2001) High level secretory production of human G-CSF by fed-batch culture of recombinant *Escherichia coli*. *Bioproc Biosystems Eng* 24:249-254
- Zavialov AV, Batchikova NV, Korpela T, Petrovskaya LE, Korobko VG, Kersley J, MacIntyre S, Zavyalov S (2001) Secretion of recombinant proteins via the chaperone/usher pathway in *Escherichia coli*. *Appl Environ Microbiol* 67:1805-1814
- Zhou S, Yomano LP, Saleh AZ, Davis FC, Aldrich HC, Ingram LO (1999) Enhancement of expression and apparent secretion of *Erwinia chrysanthemi* endoglucanase (encoded by *celZ*) in *Escherichia coli* B. *Appl Environ Microbiol* 65:2439-2445